Stenknäck (Coccothraustes coccothraustes) sågs i två ex. vid Johannisberg, V Västerås, den 7.4.1950. De höllo till högt uppe i ekar.

SVEN-OLOF ANDERSSON, HELGE KJELLSTRÖM

Kungsörn (Aquila chrysaëtos). Den 24.3.1950 sågs ett ex. segelflyga över Offerlund i Lundby s:n invid Västerås.

Sven-Olof Andersson, Helge Kjellström

Summaries:

- 1. Some observations of the Two-barred Crossbill (Loxia leucoptera). In 1929--1930, when crossbills were numerous in South Sweden, a flock of ab. 12 Loxia leucoptera was seen at Limhamn near Malmö, S. Sweden. In April, 1930, 4 specimens were observed at Skållerud, prov. Dalsland, and in the winter 1949-50 a flock of 7 birds was noted at Fröseke, prov. Småland.
- Waxwings (Bombycilla garrulus) in Gothenburg (Göteborg). Large flocks were noted in November and the beginning of December 1952. As seen from the photo. the birds alight by preference very high up in places commanding free sight in different
- 3. (a) The Little Ringed Plover (Charadrius dubius) was found breeding at lake Mollsjön, southern Västergötland in 1948 and 1952 but not in the interjacent years, owing to higher water level.
- (b) The Great Black-backed Gull (Larus marinus) has been breeding at this same lake since 1948. At the lake Stora Björsjön in the same Province two pairs were breeding in 1952 and probably some years before. This species usually does not breed at freshwater in Sweden, but is regular in lake Vänern.
- 4. The Little Ringed Plover (Charadrius dubius) breeding at Uddevalla. A nest of Charadrius dubius was found on May 21, 1952, near a railway track in the harbour of Uddevalla, prov. Bohuslan.
- 5. Nesting records of Dunlin (Calidris alpina) and Long-tailed Skua (Stercorarius longicaudus) from Härjedalen. One nest of each (and also vagrant flocks of the latter species) in the border district between Harjedalen and Jämtland in 1950. Earlier information on the breeding of these birds in Härjedalen is very scanty.
- 6. The Black-headed Gull (Larus ridibundus) in Bohuslän. Two new breeding-places are reported from the south of Bohuslan. The species now exists in scattered localities in the coastal region of this province north to Strömstad,
- 7. The effect on birds of the cold spell in the spring of 1952. This cold spell, in late March and early April following a period of favourable weather that had stimulated the migration of many birds, had a regrettable effect especially on starlings, lapwings, and certain water-fowl. Examples are given from Småland and from the island of Öland, where many birds were found dead, and others, e. g. flocks of lapwings (Vanellus vanellus), invaded the farms to be fed by man.

87315

8. WI from the rus stella (Falco ri which us first reco Feb. 18, a few da

> 9. An by crows and, acc a neighb character northerly navia di

10. R in the

Ödskölt

Fogdö Huddin Stoc Huddin Västeri Billing Billing Billing

Form from the 1943 (see southern clusively however,

11, 8 at Hall in south 28, 1952 holm, 8 dead AL